



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Following a rumor that the "California Quail" had been introduced I made inquiry of several of the settlers but found nothing to confirm such report.

The Partridge seems first to have appeared on Upper Willow Creek about 10 or 12 years ago and gradually became somewhat common. The hard winter of 1905-06, drove many flocks to the barnyards for food, where it would seem they met with almost universal destruction at the hands of settlers.

For several years they were not seen at all, but are again becoming somewhat common. No specimens have been examined, I am therefore unable to state how the species compares with those from the Cascades and Coast Mts.

The past May a small colony of two or three pairs of Bobolinks were found nesting in a meadow at Ironside, the first record I think for the state, and extending the known range of the species considerably to the westward.—A. W. ANTHONY, *Portland, Ore.*

Additions to Birds of Kerrville, Texas.—In my paper on the Birds of Kerrville, published in 'The Auk' 1911, pp. 200-219, I accidentally omitted the following species.

Charitonetta albeola. BUFFLE-HEAD.—Occasional on the Guadalupe river. A female was shot last December in the neighborhood of Kerrville.

Bartramia longicauda. UPLAND PLOVER.—Formerly very numerous during migration in the more open parts of the country, but rapidly becoming scarcer.

Aquila chrysaetos. GOLDEN EAGLE.—A young bird taken from a nest near the head waters of the Guadalupe, was kept for several years as a pet in a saloon at Kerrville. Have seen the bird two or three times on Turtle Creek. One was shot near the head of the Guadalupe river some time during last winter and the stuffed skin is in Kerrville.

Haliaeetus l. leucocephalus. BALD EAGLE.—Formerly not uncommon and used to breed on bluffs along the Guadalupe, Frio and Medina rivers. Both of these Eagles are occasionally troublesome to the ranch owners, killing young lambs and kids.

Last year I also added two species to my list of birds of the neighborhood of Kerrville.

Pelecanus erythrorhynchos. WHITE PELICAN.—On April 23, a flock of about eighty were seen along the river about three miles from Kerrville, and one of them was killed and I saw the remains.

Wilsonia pusilla pileolata. PILEOLATED WARBLER.—One male at the ranch on Turtle Creek May 11, 1911.—HOWARD LACEY, *Kerrville, Texas.*

Recent Records from the Valley of the Lower Rio Grande.—Two examples of the Yellow-crowned Night Heron (*Nyctanassa violacea*) ♀ Jan.